Impact of the COVID-19 pandemic on the detection and management of colorectal cancer in England

Updated analyses to include referrals, colonoscopies and surgical treatments up to August 2021

Introduction

Researchers in the Nuffield Department of Population Health, working with Cancer Research UK, NHS Digital, Public Health England and a team of experts from other UK universities, analysed patterns of diagnosis and care for colorectal cancer during the COVID-19 pandemic. Analyses quantified the monthly number of two-week wait referrals, colonoscopies and 31-day referrals for first treatment as well as patterns of use of surgery and rectal radiotherapy. Results based on data from NHS Hospital Trusts in England up to 31st October 2020 were published in Lancet Gastroenterology and Hepatology on the 15th January 2021¹. Updated monthly analyses are published here https://www.ndph.ox.ac.uk/corectr/covid19

Summary

There was a significant reduction in two week wait referrals in April 2020 but they had returned to 2019 levels by September 2020 and were maintained up to February 2021 with further increase up to August. The number of referrals seen within the two week target increased slightly to be above 2019 levels in November and December 2020 but fell in January before recovering slightly in February and March and then declining again up to August 2021. Colonoscopy activity returned to a level similar to 2019 by October 2020 but fell again in December before beginning to recover in subsequent months. The number of people referred for their first treatment within 31-days was above that from an average month in 2019 from October 2020 onwards but the proportion meeting the target has fallen below previous levels throughout 2021 so far.

Colorectal cancer surgery patterns in 2021 had recovered to levels similar to 2019. It was not possible to update the rectal radiotherapy analyses to include figures beyond October 2020 as Public Health England would not provide the relevant extract of the National Radiotherapy Dataset.

Updated Figures

Figure 1: Monthly number of referrals into the 2-week wait pathway and the proportion meeting 2-week target in England

Figure 2: Monthly number of colonoscopies undertaken in England

Figure 3: Monthly number of 31-day to treatment pathway referrals and the proportion of referrals meeting that target in England

Figure 4: Monthly number of operations undertaken for colorectal, colonic, and rectal cancer in England

Figure 5: Monthly number and proportion of operations for colorectal, colonic and rectal cancer that were undertaken laparoscopically


Figure 6: Monthly number and proportion of operations for colorectal, colonic and rectal cancer in which a stoma was formed

Figure 7: Monthly number and proportion of operations for colorectal, colonic and rectal cancer that were undertaken following an emergency admission

Reference

1. Morris EJA, Goldacre R, Spata E, et al. Impact of the COVID-19 pandemic on the detection and management of colorectal cancer in England: a population-based study. *Lancet Gastroenterol Hepatol* 2021; **6**(3): 199-208.

Figure 1: Monthly number of referrals into the 2-week wait pathway and the proportion meeting 2-week target in England



Error bars represent +/-1 SD of the (pre-COVID-19) monthly counts for 2019.

Figure 2: Monthly number of colonoscopies undertaken in England

Error bars represent +/- 1 SD of the (pre-COVID-19) monthly counts for 2019





Figure 3: Monthly number of 31-day to treatment pathway referrals and the proportion of referrals meeting that target in England



Error bars represent +/-1 SD of the (pre-COVID-19) monthly counts for 2019.

Figure 4: Monthly number of operations undertaken for colorectal, colonic, and rectal cancer in England

Error bars represent +/- 1 SD of the (pre-COVID-19) monthly counts for 2019. The 2021 rate is based on counts adjusted for incomplete coding




Figure 5: Monthly number and proportion of operations for colorectal, colonic and rectal cancer that were undertaken laparoscopically

Error bars represent +/- 1 SD of the (pre-COVID-19) monthly counts for 2019. The 2021 rate is based on counts adjusted for incomplete coding




Figure 6: Monthly number and proportion of operations for colorectal, colonic and rectal cancer in which a stoma was formed

Error bars represent +/- 1 SD of the (pre-COVID-19) monthly counts for 2019. The 2021 rate is based on counts adjusted for incomplete coding




Figure 7: Monthly number and proportion of operations for colorectal, colonic and rectal cancer that were undertaken following an emergency admission

Error bars represent +/- 1 SD of the (pre-COVID-19) monthly counts for 2019. The 2021 rate is based on counts adjusted for incomplete coding.

